



ADDVET

PRODUCTS CATALOG

Antibiotic &
Anticoccidial

addvet.net



ADDVET

PRODUCTS CATALOG

Antibiotic & Anticoccidial



addvet.net

COMPANY PROFILE

Who we are

Established in 2014 with a capital of one million dollars, ADDVET is a Jordanian company specialized in manufacturing veterinary drugs.

What we offer

A variety of veterinary medicines Powders & Liquids orals, including Antibiotics, Vitamins, Amino Acids, Anticoccidial drugs & Premixes.

Our Mission

A variety of veterinary medicines Powders & Liquids orals, including Antibiotics, Vitamins, Amino Acids, Anticoccidial drugs & Premixes.

Our Vision

To improve the world feed and food business and be part of the next generation global feed chain.



addvet.net



ANTIBIOTIC

ADDCIPRO
ADDENRO
ADDFLOR
ADDFOS
ADDTILMO
ENROHEXIN
TYFOS
TYLOHEXIN
ADDAMOX
ADDDOX
ADDOXY
ADDTHRO

AMOXYCOL
AMPICOL
DOXYCOL
GENTDOXY
GENTTYLO
TYLODOXY
TYLOTHRO
ADDAMPI-50
ADDCOL12
TYLOFORT



ADDCIPRO

Acts bactericidally to treat gastrointestinal, respiratory and urinary tract infections.

Composition

Each 1 ml contains:
Ciprofloxacin 100 mg

Ciprofloxacin belongs to the group of quinolones and acts bactericidal against mainly Gramnegative bacteria like Campylobacter, E. coli, Haemophilus, Mycoplasma, Pasteurella and Salmonella spp.

Indications

To treat ciprofloxacin sensitive microorganisms in poultry, calves, goat and sheep that cause gastrointestinal, respiratory and urinary tract infections.

Dosage and Administration

Poultry: 1 Litre / 1000 Litres of drinking water
or as advised by a veterinarian

Calves, goats and sheep:

Twice daily 10 ml per 75 - 150 kg body weight for 3 - 5 days.

Withdrawal times

- For meat: 12 days.



ADDENRO

Acts bactericidally to treat gastrointestinal, respiratory and urinary tract infections.

Composition

Each 1 ml contains:

Enrofloxacin 200 mg

Enrofloxacin belongs to the group of quinolones and acts bactericidal against mainly Gram-negative bacteria like Campylobacter, E. coli, Haemophilus, Pasteurella, Salmonella and Mycoplasma spp.

Indications

To treat enrofloxacin sensitive microorganisms in poultry, calves, goat and sheep that cause gastrointestinal, respiratory and urinary tract infections.

Dosage and Administration

Poultry: 500 ml / 1000 Litres of drinking water or as advised by a veterinarian.

Calves, goats and sheep:

Twice daily 5 ml per 75 - 150 kg body weight for 3 - 5 days.

Withdrawal times

- For meat and offal:

Chickens : 7 days.

Turkeys : 13 days.



KEEP OUT OF REACH OF CHILDREN.

STORE PRODUCT IN A COOL PLACE.

4

ADDFLOR

A bacteriostatic to treat gastrointestinal, and respiratory tract infections in animals.

Composition

Each 1 ml contains:

Florfenicol 100 mg

Florfenicol is a synthetic broad-spectrum antibiotic that is effective against most Gram-positive and Gram-negative bacteria isolated from domestic animals. Florfenicol, a fluorinated derivative of chloramphenicol acts by inhibiting protein synthesis at the ribosomal level and is bacteriostatic.

Indications

Preventive and therapeutic treatment of gastrointestinal and respiratory tract infections, caused by florfenicol sensitive microorganisms such as Actinobaccillus spp., Pasteurella spp., Salmonella spp. and Streptococcus spp. in poultry.

Dosage and Administration

Poultry: 1 Litre / 1000 Litres of drinking water or as advised by a veterinarian

Swine: 1 litre per 1000 litres of drinking water (100 ppm; 10 mg/kg or 10 ml per 100 kg) for 5 days.

Withdrawal times

- For meat: Poultry : 7 days.



FOR VETERINARY USE ONLY.

5

ADDFOS

Broad-spectrum bactericidal antimicrobial of fosfonic acid family to treat diseases.

Composition

Each 1 ml contains:

Fosfomycin (as sodium) 100 mg

Fosfomycin is a broad-spectrum bactericidal antimicrobial belonging to the fosfonic acid family, recommended as an aid for the treatment of bacterial diseases of poultry.

Indications

For the treatment of bacterial infections of poultry susceptible to the action of fosfomycin such as Infectious coryza, fowl cholera, colibacillosis, ORT, CRD, staphylococcal infections, fowl thyphoid, and salmonellosis.

Dosage and Administration

Poultry : 500 ml / 1000 liters of drinking water for 3-5 consecutive days.

Calves, goats and sheep :

Twice daily 10 ml per 75 - 150 kg body weight for 3 - 5 days.

Withdrawal times

- For meat: Poultry : 3 days.



KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

6

ADDTILMO

Broad-spectrum semi-synthetic bactericidal macrolide antibiotic to treat respiratory diseases in poultry.

Composition

Each 1 ml contains:

Tilmicosin phosphate 250 mg

Tilmicosin is a macrolide antibiotic. It is used in veterinary medicine for the treatment of bovine respiratory disease and bovine respiratory disease associated with Mannheimia (Pasteurella) haemolytica.

Indications

For the control and treatment of respiratory infections associated with tilmicosin-susceptible micro-organisms such as Mycoplasma spp. Pasteurella multocida, Actinobacillus pleuropneumoniae, Actinomyces pyogenes and Mannheimia haemolytica in chickens and turkeys.

Dosage and Administration

For oral administration.

Calves:

Twice daily, 1 ml per 20 kg body weight via (artificial) milk for 3 - 5 days.

Poultry:

300 ml per 1000 litres of drinking water for 3 days.

Withdrawal times

- For meat:

Calves : 42 days.

Broilers : 12 days.

Turkeys : 19 days.



FOR VETERINARY USE ONLY.

7

ENROHEXIN

Quinolone antibiotic with added bromhexine to treat respiratory tract diseases in animals.

Composition

Each 1 ml contains:

Enrofloxacin 100 mg Bromhexine 10 mg

Enrofloxacin belongs to the group of quinolones and acts bactericidal against mainly Gram-negative bacteria like Campylobacter, E. coli, Haemophilus, Pasteurella, Salmonella and Mycoplasma spp. Bromhexine is a mucolytic.

Indications

For the treatment and control of respiratory infections associated with Mycoplasma spp., Pasteurella multocida., Haemophilus paragalinarum, Chronic Respiratory Disease (CRD), in some infections produced by susceptible strains of Staphylococcus, and also for other enrofloxacin-susceptible bacteria.

Dosage and Administration

Poultry: 1 Litres / 1000 Litres of drinking water or as advised by a veterinarian.

Calves, goats and sheep:

Twice daily 5 ml per 75 - 150 kg body weight for 3 - 5 days.

Withdrawal times

- For meat: 12 days.



KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

8

TYFOS

Bacteriostatic action against mainly Gram-positive and Gram-negative bacteria.

Composition

Each 1 ml contains:

Tylosin 200 mg
Fosfomicin 100 mg

Indications

It is a combination of broad-spectrum antibiotics indicated for the treatment of poultry diseases, caused by Gram-positive, Gram-negative and Mycoplasmas. Indicated in cases of mycoplasmosis, colibacillosis, chronic respiratory disease in poultry, Staphylococcus infections, Listeriosis, Avian Cholera, infectious Coryza, and diseases caused by Fosfomicin and/or Tylosin susceptible organisms.

Dosage and Administration

For oral administration via drinking water.

Calves : 4 ml per 50 kg bodyweight twice daily, during 7 – 14 days.

Poultry : 250 ml per 100 litres of drinking water during 3 – 5 days.

Withdrawal times

Meat : 15 days

Eggs : 4 days



FOR VETERINARY USE ONLY.

9

TYLOHEXIN

Macrolide antibiotic with added bromhexine to treat respiratory tract diseases.

Composition

Each 1 ml contains:

Tylosin 200 mg
Bromhexine 10 mg

Indications

For the treatment and control of respiratory infections associated with Mycoplasma spp., Pasteurella multocida., Haemophilus paragalinarum, Chronic Respiratory Disease (CRD), in some infections produced by susceptible strains of Staphylococcus, and also for other Tylosin-susceptible bacteria.

Dosage and Administration

For oral administration via drinking water.

Calves : 4 ml per 50 kg bodyweight twice daily, during 7 – 14 days.

Poultry : 500-1000 ml per 1000 litres of advised by a veterinarian.

Withdrawal times

For meat:

Calves, goats and sheep : 14 days.

Poultry : 7 days.



KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

10

ADDAMOX

Broad-spectrum penicillin with a bactericidal action against both Gram+ve and Gram-ve bacteria.

Composition

Each 1 gm contains:

Amoxicillin 300 mg

Indications

Gastrointestinal, respiratory, and urinary tract infections caused by amoxicillin sensitive micro-organisms, like Campylobacter, Clostridium, Corynebacterium, E. coli, Erysipelothrix, Haemophilus, Pasteurella, Salmonella, penicillinase negative Staphylococcus and Streptococcus spp. in animals.

Dosage and Administration

Poultry: 500 gm / 1000 Litres of drinking water

or as advised by a veterinarian.

Calves : 10 g per 100 kg bodyweight daily, during 3 – 5 days.

Withdrawal times

Meat : 7 days

Milk : 3 days

Not for use in laying birds producing eggs for human consumption



FOR VETERINARY USE ONLY.

11

ADDDOX

Acts bacteriostatic against many Gram-positive and Gram-negative bacteria in animals.

Composition

Each 1 gm contains:
Doxycycline hyclate 500 mg

Doxycycline belongs to the group of tetracyclines and acts bacteriostatic against many Gram-positive and Gram-negative bacteria. The action of doxycycline is based on the inhibition of bacterial protein synthesis.

Indications

Gastrointestinal and respiratory infections caused by doxycycline sensitive micro-organisms, like Bordetella, Campylobacter, Chlamydia, E. coli, Haemophilus, Mycoplasma, Pasteurella, Rickettsia, Salmonella, Staphylococcus, and Streptococcus spp. in animals.

Dosage and Administration

For oral administration via drinking water (poultry) or milk replacer (calves).
Calves : 1 g per 50 kg bodyweight, once daily during 3 – 5 days.
Poultry :
100 – 250 g per 1,000 litres of drinking water during 3 – 5 days.
Medicated drinking water should be used within 24 hours.
Medicated milk replacer should be used within 12 hours.

Withdrawal times

- For meat:
Calves, goats and sheep : 14 days.
Poultry : 7 days.



KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

12

ADDOXY

Broad-spectrum tetracycline with a bactericidal action against both Gram+ve and Gram-ve bacteria.

Composition

Each 1 gm contains:
Oxytetracycline 500 mg

The mode of action of oxytetracycline is based on the inhibition of bacterial protein synthesis.

Indications

Gastrointestinal and respiratory infections caused by oxytetracycline sensitive bacteria like Bordetella, Bacillus, Corynebacterium, Campylobacter, E. coli, Haemophilus, Pasteurella, Salmonella, Staphylococcus and Streptococcus spp. and Mycoplasma, Rickettsia and Chlamydia spp. in various animals.

Dosage and Administration

For oral administration via feed or drinking water or milk replacer.
Poultry:
500 g per 1000 – 1500 litres of drinking water during 3 – 5 days.
Calves, sheep, goats, lambs, kids : 80 mg per kg bodyweight daily, during 3 – 5 days.

Withdrawal times

Meat :
Poultry : 7 days
Other : 8 days
Eggs : 5 days



FOR VETERINARY USE ONLY.

13

ADDTHRO

Reliable for Chronic Respiratory Disease (CRD) caused by Mycoplasma infections in poultry.

Composition

Each 1 gm contains:
Erythromycin thiocyanate 350 mg

Erythromycin binds reversibly to the 50S subunit of ribosomes, resulting in the blockage of transpeptidation or translocation reactions, leading to impaired and/or inhibited protein synthesis and to inhibit cell growth.

Indications

Chronic Respiratory Disease (CRD) is caused by Mycoplasma infections, and for the treatment of infectious respiratory diseases due to erythromycin-sensitive bacteria such as Staphylococcus aureus (e.g. infectious arthritis), Haemophilus paragallinarum (e.g. infectious coryza) and Pasteurella multocida (e.g. pasteurellosis; pneumonia) in poultry.

Dosage and Administration

Poultry: 500 gm / 1000 Litres of drinking water or as advised by a veterinarian

Withdrawal times

- For meat : 3 days.
- For eggs : 6 days.



KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

14

AMOXYCOL

Quality duo medicinal combination with bactericidal action against many Gram+ve and Gram-ve bacteria.

Composition

Each 1 gm contains:
Amoxicillin trihydrate 200 mg
Colistin sulphate 1200000 I.U.

Indications

Gastrointestinal, respiratory, and urinary tract infections caused by amoxicillin and colistin sensitive micro-organisms, like Campylobacter, Clostridium, Corynebacterium, E. coli, Erysipelothrix, Haemophilus, Pasteurella, Salmonella, penicillinase negative Staphylococcus and Streptococcus spp. in animals.

The action of amoxicillin is based on the inhibition of cell wall synthesis. Colistin interacts strongly with phospholipids and penetrates and disrupt the structure of cell membranes.

Dosage and Administration

Poultry: 500 gm / 1000 Litres of drinking water or as advised by a veterinarian.
Calves : 10 g per 75 kg bodyweight daily, during 3 – 5 days.

Withdrawal times

- Meat : 7 days
- Milk : 3 days
- Eggs : 3 days



FOR VETERINARY USE ONLY.

15

AMPICOL

Quality duo combination with bactericidal action against many Gram+ve and Gram-ve bacteria.

Composition

Each 1 gm contains:

Ampicillin trihydrate 200 mg
Colistin sulphate 1200000 I.U.

Indications

Gastrointestinal and urinary tract infections caused by ampicillin and colistin sensitive micro-organisms, like Campylobacter, Clostridium, Corynebacterium, E. coli, Erysipelothrix, Haemophilus, Pasteurella, Salmonella, penicillinase negative Staphylococcus, and Streptococcus spp. in animals. It is effective for diarrhea and toxicity cases.

Dosage and Administration

Poultry: 500 gm / 1000 Litres of drinking water
or as advised by a veterinarian.

Calves : 10 g per 75 kg bodyweight daily, during 3 – 5 days.

Withdrawal times

Meat : Calves, Poultry : 2 days



KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

16

DOXYCOL

Quality duo medicinal combination acting bacteriostatic against many Gram+ve and Gram-ve bacteria.

Composition

Each 1 gm contains:

Doxycycline Hyclate 200 mg
Colistin sulphate 1200000 I.U.

Indications

Gastrointestinal and respiratory infections (GRD) are caused by microorganisms sensitive to colistin and/or doxycycline like Bordetella, Campylobacter, Chlamydia, E. coli, Klebsiella, Haemophilus, Mycoplasma, Pasteurella, Rickettsia, Salmonella, Staphylococcus and Streptococcus spp. in animals.

The action of doxycycline is based on the inhibition of bacterial protein synthesis. Colistin interacts strongly with phospholipids and penetrates It can cause cell membranes to break down.

Dosage and Administration

Poultry: 500 gm / 1000 Litres of drinking water or
as advised by a veterinarian.

Calves, goats and sheep:

Twice daily 5 g per 200 kg body weight for 3 - 5 days.

Withdrawal times

- For meat:

Calves, goats and sheep : 14 days.

Poultry : 7 days.



FOR VETERINARY USE ONLY.

17

GENTDOXY

Duo antibiotic class that acts bacteriostatic against many Gram+ve and Gram-ve bacteria.

Composition

Each 1 gm contains:

Doxycycline Hyclate	200 mg
Gentamicin sulphate	200 mg

The action of doxycycline and gentamicin is based on the inhibition of bacterial protein synthesis.

Indications

Gastrointestinal and respiratory infections caused by microorganisms sensitive to doxycycline and/or gentamicin like Bordetella, Campylobacter, Chlamydia, E. coli, Klebsiella, Haemophilus, Mycoplasma, Pasteurella, Rickettsia, Salmonella, Staphylococcus and Streptococcus spp. in animals.

Dosage and Administration

Poultry: 500 gm / 1000 Litres of drinking water or as advised by a veterinarian.

Calves, goats and sheep:

Twice daily 5 g per 200 kg body weight for 3 - 5 days.

Withdrawal times

- For meat:

Calves, goats and sheep : 14 days.

Poultry : 7 days.



**KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.**

18

GENTTYLO

Duo antibiotic class that acts bacteriostatic against many Gram+ve and Gram-ve bacteria.

Composition

Each 1 gm contains:

Tylosin tartrate	200 mg
Gentamicin sulphate	200 mg

The action of both tylosin and gentamicin is based on the inhibition of bacterial protein synthesis.

Indications

Gastrointestinal and respiratory infections caused by microorganisms sensitive to tylosin and/or gentamicin like Bordetella, Campylobacter, Chlamydia, E. coli, Klebsiella, Haemophilus, Mycoplasma, Pasteurella, Rickettsia, Salmonella, Staphylococcus and Streptococcus spp. in animals.

Dosage and Administration

Poultry: 500 gm / 1000 Litres of drinking water or as advised by a veterinarian.

Calves, goats and sheep:

Twice daily, 5 g per 100 kg body weight for 3 - 5 days.

Withdrawal times

Meat : 15 days

Eggs : 4 days



FOR VETERINARY USE ONLY.

19

TYLODOXY

Quality duo medicinal combination acting bacteriostatic against many Gram+ve and Gram-ve bacteria.

Composition

Each 1 gm contains:

Tylosin tartrate	120 mg
Doxycycline Hyclate	200 mg

The action of tylosin and doxycycline is based on the inhibition of bacterial protein synthesis. They bind reversibly to the 50S subunit of ribosomes.

Indications

Gastrointestinal and respiratory infections (CRD) are caused by microorganisms sensitive to tylosin and/or doxycycline like Bordetella, Campylobacter, Chlamydia, E. coli, Klebsiella, Haemophilus, Mycoplasma, Pasteurella, Rickettsia, Salmonella, Staphylococcus and Streptococcus spp. in animals.

Dosage and Administration

Poultry: 500 gm / 1000 Litres of drinking water or as advised by a veterinarian.

Calves, goats and sheep:

Twice daily 5 g per 200 kg body weight for 3 - 5 days.

Withdrawal times

- For meat:

Calves, goats, poultry and sheep : 5 days.



KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

20

TYLOTHRO

Quality duo macrolide class acting bacteriostatic against many Gram+ve and Gram-ve bacteria.

Composition

Each 1 gm contains:

Tylosin tartrate	120 mg
Erythromycin thiocyanate	200 mg

The action of tylosin and erythromycin is based on the inhibition of bacterial protein synthesis. They bind reversibly to the 50S subunit of ribosomes.

Indications

Gastrointestinal and respiratory infections (CRD) are caused by microorganism sensitive to tylosin and/or erythromycin like Bordetella, Campylobacter, Chlamydia, E. coli, Klebsiella, Haemophilus, Mycoplasma, Pasteurella, Rickettsia, Salmonella, Staphylococcus and Streptococcus spp. in animals.

Dosage and Administration

Poultry: 500 gm / 1000 Litres of drinking water or as advised by a veterinarian.

Calves, goats and sheep:

Twice daily, 5 g per 100 kg body weight for 3 - 5 days.

Withdrawal times

- For meat:

Calves, goats, poultry and sheep : 5 days.



FOR VETERINARY USE ONLY.

21

ADDAMPI-50

Quality duo to acts as bactericidal action against many Gram+ve and Gram-ve bacteria.

Composition

Each 1 gm contains:
Ampicillin (Trihydrate) 500 mg

Indications

Ampicillin is indicated for the treatment of infections caused by E.coli, Clostridium, Salmonella, Brucella, Proteus, Klebsiella, Strepto and Staphylococci, Corynebacterium, Haemophilus, Pasteurella, and Shigella in the different animals. Ampicillin 50% is recommended for use in cases of Bacterial enteritis and Necrotic enteritis in poultry. It is also indicated Against Pneumonia, Septicemia, Diarrhea, and Scours in calves and lambs.

Dosage and Administration

Poultry: 100 gm / 500 Litres of drinking water
or as advised by a veterinarian.

Calves, lambs, kids :

10 -20 g per 200 kg bodyweight, twice daily, during 5 days.

Withdrawal times

Meat : Calves : 2 days

Poultry : 1 day



KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

22

ADDCOL12

Acts bacteriostatic against many Gram-positive and Gram-negative bacteria in animals.

Composition

Each 1 gm contains:
Colistin sulphate 1200000 IU.

Indications

ADDCOL12 Powder is indicated for the treatment of enteric infections caused by gram-negative bacteria e.g. E. coli, Salmonella, and Shigella in poultry, calves, and lambs. Colistin is highly effective against strains of E. coli, Pseudomonas aeruginosa, Salmonella spp., Haemophilus spp., Shigella spp., Pasteurella spp., Brucella spp., Aerobacteraerogenes, and Bordetella bronchiseptica. Gram-positive bacteria are generally less sensitive. However, there are sensitive strains of Staphylococcus spp., Bacillus spp., Streptococcus pyogenes and Corynebacterium spp.

Dosage and Administration

Calves, goats, and sheep:

Twice daily 5 g per 100 kg body weight for 5 - 7 days.

Poultry: 500 gm per 500 - 1000 liters of drinking water
or 250 - 500 kg of feed for 5 - 7 days.

Withdrawal times

Meat : 3 days

Eggs : 5 days



FOR VETERINARY USE ONLY.

23

TYLOFORT

Acts bacteriostatic against many Gram-positive and Gram-negative bacteria in animals.

Composition

Each 500 gm contains:

Tylosin active (Tartrate) 500 gm

Tylosin antibiotics belong to macrolide group and it's highly effective against Mycoplasma, Gram-positive bacteria and is absorbed rapidly from the GIT.

Indications

TYLOFORT is highly effective against Mycoplasma, Chronic respiratory disease (CRD) and gram-positive bacteria.

TYLOFORT used for the prevention and treatment of many respiratory tract diseases such as; Air sacculitis, Infectious synovitis, Infectious sinusitis, Infectious coryza, Spirochaetosis, Chlamydia, and stress conditions that affect Poultry.

Dosage and Administration

Poultry: Prevention: 50 gm / 200 Liter of drinking water for 3-5 days.

Treatment: 100 gm / 200 Liter of drinking water for 3-5 days.

Calves, goats and sheep:

Twice daily 5 g per 220 - 250 kg body weight for 5 - 7 days.

Withdrawal times

Meat: Calves : 7 days Poultry : 5 days

Eggs : 4 days



ANTICOCCIDIAL

ADDAMPRO

ADDTOLCOX

AMPROCOX-K3



KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

24

25



ADDAMPRO

A reliable anticoccidial and antiprotozoal agent to treat coccidiosis in animals.

Composition

Each 1 ml contains:
Amprolium HCL 600 mg

Indications

Amprolium is an anticoccidial with activity against Eimeria spp., especially Eimeria tenella and Eimeria necatrix. Helps to treat coccidiosis in calves, sheep, goats, and poultry. Coccidiosis of all stages like schizogony and gametogony stages of Eimeria spp. in animals. Amprolium is a coccidiostat and antiprotozoal agent used for the treatment and prevention of coccidiosis. The anticoccidial effect of Amprolium is achieved by inhibiting the Eimeria spp. to uptake thiamine.

Dosage and Administration

Poultry:
250 ml per 1000 litres of drinking water during 5 days.
Treatment: 500 ml per 1000 litre of drinking water during 5-7 days.

Withdrawal times

- For meat:
Calves, goats, sheep : 3 days.
Poultry : 3 days.



KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

26

ADDTOLCOX

A reliable anticoccidial with activity against Eimeria spp. especially Eimeria tenella and Eimeria necatrix in animals.

Composition

Each 1 ml contains:
Toltrazuril 25 mg

Indications

Toltrazuril is an anticoccidial with activity against Eimeria spp., especially Eimeria tenella and Eimeria necatrix. Helps to treat coccidiosis in calves, sheep, goats, and poultry. Coccidiosis of all stages like schizogony and gametogony stages of Eimeria spp. in animals.

Dosage and Administration

For oral administration:
500 ml per 500 Litres of drinking water for continuous medication over 48 hours or 1500 ml per 500 Litres of drinking water given for 8 hours per day, on 2 consecutive days. This corresponds to a dose rate of 7 mg of toltrazuril per kg of body weight per day for 2 consecutive days.

Withdrawal times

For meat:
- Chickens : 18 days.
- Turkeys : 21 days.



FOR VETERINARY USE ONLY.

27

AMPROCOX-K3

A reliable anticoccidial with activity against Eimeria spp. especially Eimeria tenella and Eimeria necatrix in animals.

Composition

Each 1 ml contains:

Amprolium HCL 250 mg

Vitamin K3 100 mg

Indications

Amprolium is an anticoccidial with activity against Eimeria spp., especially Eimeria tenella and Eimeria necatrix. Helps to treat coccidiosis in calves, sheep, goats, and poultry. Coccidiosis of all stages like schizogony and gametogony stages of Eimeria spp. in animals.

Dosage and Administration

For oral administration:

500 ml per 500 Litres of drinking water for continuous medication over 48 hours or 1500 ml per 500 Litres of drinking water given for 8 hours per day, on 2 consecutive days.

Withdrawal times

- For meat:

Calves, goats, sheep : 3 days.

Poultry : 3 days.



KEEP OUT OF REACH OF CHILDREN.

STORE PRODUCT IN A COOL PLACE.

28

ANTHELMINTIC

BENDA-M

LEVACTIZOLE 6 IN 1



ADDvet
Animal Health

29

BENDA -M

Benda-M is a broad spectrum multi-purpose anthelmintic for the control of mature and developing immature forms of gastrointestinal roundworms, lungworms, tapeworms and adult liver fluke in cattle, camels, sheep & goats. The product is also ovicidal against fluke and roundworm eggs. In sheep it is active against benzimidazole-susceptible strains

Composition

Each 1 ml of Benda-M contains:

Albendazole	25mg
Selenium (as sodium selenite)	0.27mg
Cobalt (as cobalt sulphate)	0.62mg

Indications

In cattle & camels it is active against the following species:

Roundworms: Ostertagia, Haemonchus, Trichostrongylus, Nematodirus, Oesophagostomum, Bunostomum, Cooperia and Strongyloides species. It is usually effective against inhibited larvae of Cooperia and Ostertagia. Lungworms: Dictyocaulus viviparus, Tapeworms: Moniezia species, Adult liver fluke: Fasciola hepatica.

In goats it is active against benzimidazole-susceptible strains of the following species: Roundworms: Ostertagia, Haemonchus, Trichostrongylus, Nematodirus (including N battus), Chabertia and Oesophagostomum. It is usually effective against inhibited larvae of Ostertagia. Lungworms: Dictyocaulus filaria. Tapeworms: Moniezia species. Adult liver fluke: Fasciola hepatica.

BENDA-M is ovicidal and will kill fluke and roundworm eggs, thus reducing pasture contamination.

The selenium and cobalt in this product are trace elements of use as nutritional supplement.

Dosage and Administration

Sheep & Goats:

Worm dose 2ml per 10kg

Fluke & amp; Worm dose 3ml per 10kg

Cattle & Camels:

Worm dose 15ml per 50kg

Fluke & amp; Worm dose 20ml per 50kg of drinking water given for 8 hours per day, on 2 consecutive days.

Withdrawal times

- For meat:

Calves, goats, sheep : 4 days.

Milk: 60 Hours



KEEP OUT OF REACH OF CHILDREN.

STORE PRODUCT IN A COOL PLACE.

30

LEVACTIZOLE 6 IN 1

LEVACTIZOLE 6 IN 1 Oral Suspension is used for the treatment of and prevention from external and internal infections caused by the parasites in cattle, camels, sheep and goats.

Composition

Each liter Contains:	Sodium selenite: 160 mg
Albendazole: 25 g	Ivermectin: 1 gm
Cobalt sulphate: 1500 mg	Zinc: 0.6 gm
Levamisole Hcl: 7.5 gm	

Indications

Gastro-Intestinal nematodes : Ostertagia sp., Haemonchus sp., Trichostrongylus sp., Cooperia., Oesophagostomum sp., Bunostomum sp and Chabertia sp., Tenia : Moniezia sp.

Pulmonary Enterobiasis : Dictyocaulus viviparus.

Hepatic Fasciola : Dictyocaulus viviparus.

Hepatic Fasciola : Fasciola hepatica

Dosage and Administration

For Cattle and Camels:

It is administered at a dose of 15 ml / 50 kg bodyweight via oral route.

For hepatic fasciola, it is administered at a dose of 20 ml / 50 kg bodyweight.

For Sheep and Goats:

It is administered at a dose of 2 ml / 10 kg bodyweight via oral route

For hepatic fasciola, it is administered at a dose of 3 ml / 10 kg bodyweight

Withdrawal times

Meat: 14 days.

Milk: 60 Hours



FOR VETERINARY USE ONLY.

31

ANTIBIOTIC



ANTICOCCIDIAL & SUSPENSION



KEEP OUT OF REACH OF CHILDREN

STORE PRODUCT IN A COOL PLACE

addvet.net



Contact

Al-Zarqa Industrial Zone [1Z4], Jordan.
info@addvet.net

